


Thyromimetic Organic Compounds

This application is a continuation in part of Application No. 09/533,219 filed March 23, 2000, which in turn claims the benefit of provisional Application No. 60/183,030 filed March 29, 1999 (converted from Application No. 09/280,105), said applications being incorporated herein by reference.

Summary of the Invention

The present invention relates to novel compounds of the formula



in which

W is O, S, S(O) or S(O)₂;

X is -SR₄, -S(O)R₄, -S(O)₂R₄, or -S(O)₂NR₅R₆; or X is -C(O)NR₅R₆ provided that -C(O)NR₅R₆ is located at the 3', 4'- or 5'-position;

Y is O or H₂;

Z is hydrogen, halogen, hydroxy, optionally substituted alkoxy, aralkoxy, acyloxy or alkoxycarbonyloxy;

R is hydrogen, halogen, trifluoromethyl, lower alkyl or cycloalkyl;

Q is 5-tetrazolyl; or Q is -C(O)R₁ wherein R₁ is hydroxy, optionally substituted alkoxy, aryloxy, heteroaryloxy, aralkoxy, cycloalkoxy, heteroaralkoxy or -NR₅R₆;

R₂ is hydrogen, halogen or alkyl;

R₃ is halogen or alkyl;

09533219 09533219 09533219

R4 is optionally substituted alkyl, aryl, aralkyl, heteroaralkyl or heteroaryl;

R5, R6 and R7 are independently hydrogen, optionally substituted alkyl, cycloalkyl, aryl, aralkyl, heteroaryl, or heteroaralkyl; or R5 and R6 combined are alkylene optionally interrupted by O, S, S(O), S(O)₂ or NR7 which together with the nitrogen atom to which they are attached form a 5- to 7- membered ring;

n represents zero or an integer from 1 to 4; and pharmaceutically acceptable salts thereof.

The compounds of the invention are thyromimetic agents which can be used to prevent and/or treat diseases associated with an imbalance of thyroid hormones, such as hypo- and hyper-thyroidism, obesity, osteoporosis and depression. The compounds of the invention are, in particular, hypolipedemic agents which enhance the clearance of cholesterol from circulation, particularly the clearance of cholesterol in the form of low density lipoproteins (LDL). They, inter alia, upregulate hepatic LDL receptor function in mammals. Thus, they are useful for reducing total cholesterol plasma levels in mammals, in particular for reducing levels of LDL-cholesterol. Furthermore, such compounds also lower elevated lipoprotein (a) [Lp(a)] levels, an independent cardiovascular risk factor, in mammals. The compounds of the invention can therefore be used for the prevention and/or treatment of occlusive cardiovascular conditions in which hyperlipidemia and hyperlipoproteinemia are implicated, such as atherosclerosis and coronary heart disease in mammals.

Detailed Description of the Invention

The present invention provides for compounds of formula I, pharmaceutical compositions comprising such compounds and for methods of using such compounds.

Listed below are definitions of various terms used to describe the compounds of the instant invention. These definitions apply to the terms as they are used throughout the specification (unless they are otherwise limited in specific instances either individually or as part of a larger group).

0096960-0969601

The term "optionally substituted alkyl" refers to unsubstituted or substituted straight or branched chain hydrocarbon groups having 1 to 20 carbon atoms, preferably 1 to 7 carbon atoms. Exemplary unsubstituted alkyl groups include methyl, ethyl, propyl, isopropyl, n-butyl, t-butyl, isobutyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl and the like. Substituted alkyl groups include, but are not limited to, alkyl groups substituted by one or more (e.g. two or three) of the following groups: halo, lower alkenyl, hydroxy, cycloalkyl, alkanoyl, alkoxy, alkyloxyalkoxy, alkanoyloxy, amino, alkylamino, dialkylamino, dialkylaminocarbonyl, alkanoylamino, thiol, alkylthio, alkylthiono, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, sulfonamido, nitro, cyano, carboxy, alkoxycarbonyl, aryl, aralkyl, aralkoxy, guanidino, heterocyclyl including indolyl, imidazolyl, furyl, thienyl, thiazolyl, pyrrolidyl, pyridyl, pyrimidyl, piperidyl, morpholinyl and the like. Preferred substituents of substituted alkyl, especially of substituted alkyl of variable R1 being substituted alkoxy, are lower alkyl, cycloalkyl, lower alkenyl, benzyl, mono or disubstituted lower alkyl, e.g. ω -(amino, mono- or di-lower alkylamino, carboxy, lower alkoxycarbonyl)-lower alkyl, α -(lower alkanoyloxy, lower alkoxycarbonyl or di-lower alkylaminocarbonyl)-lower alkyl, such as pivaloyloxy-methyl.

The term "lower alkyl" refers to those alkyl groups as described above having 1 to 7, preferably 1 to 4 carbon atoms.

The term "halogen" or "halo" refers to fluorine, chlorine, bromine and iodine.

The term "alkenyl" refers to any of the above alkyl groups having at least 2 carbon atoms and further containing at least one carbon to carbon double bond. Groups having two to four carbon atoms are preferred.

The term "alkylene" refers to a straight chain bridge of 1 to 6 carbon atoms connected by single bonds (e.g., $-(CH_2)_x-$ wherein x is 1 to 6), which may be substituted with 1 to 3 lower alkyl groups.

The term "cycloalkyl" refers to cyclic hydrocarbon groups of 3 to 8 carbon atoms.

The term "alkoxy" refers to alkyl-O-.

096660-096660

The term “acyl” refers to alkanoyl, aroyl, heteroaroyl, arylalkanoyl or heteroarylalkanoyl.

The term "alkanoyl" refers to alkyl-C(O)-.

The term "alkanoyloxy" refers to alkyl-C(O)-O-.

The terms "alkylamino" and "dialkylamino" refer to (alkyl)NH- and (alkyl)₂N-, respectively.

The term "alkanoylamino" refers to alkyl-C(O)-NH-.

The term "alkylthio" refers to alkyl-S-.

The term "alkylthiono" refers to alkyl-S(O)-.

The term "alkylsulfonyl" refers to alkyl-S(O)₂-.

The term "alkoxycarbonyl" refers to alkyl-O-C(O)-.

The term "alkoxycarbonyloxy" refers to alkyl-O-C(O)O-.

The term "alkyl" as referred to in the above definitions relates to optionally substituted alkyl as defined above.

The term "aryl" refers to monocyclic or bicyclic aromatic hydrocarbon groups having 6 to 12 carbon atoms in the ring portion, such as phenyl, naphthyl, tetrahydronaphthyl, and biphenyl groups, each of which may optionally be substituted by one to four substituents such as alkyl, halo, hydroxy, alkoxy, alkanoyl, alkanoyloxy, amino, alkylamino, dialkylamino, alkanoyl-amino, thiol, alkylthio, nitro, cyano, carboxy, carboxyalkyl, alkoxycarbonyl, alkyl-thiono, alkyl-sulfonyl, sulfonamido, heterocyclyl and the like.

The term “monocyclic aryl” refers to optionally substituted phenyl as described under aryl.

The term "aralkyl" refers to an aryl group bonded directly through an alkyl group, such as benzyl.

The term "aralkoxy" refers to an aryl group bonded through an alkoxy group.

The term "arylsulfonyl" refers to aryl-S(O)₂-.

The term "aroyl" refers to aryl-C(O)-.

The term "heterocyclyl" refers to an optionally substituted, fully saturated or unsaturated, aromatic or nonaromatic cyclic group, for example, which is a 4 to 7 membered monocyclic, 7 to 11 membered bicyclic, or 10 to 15 membered tricyclic ring system, which has at least one heteroatom in at least one carbon atom-containing ring. Each ring of the heterocyclic group containing a heteroatom may have 1, 2, 3 or 4 heteroatoms selected from nitrogen atoms, oxygen atoms and sulfur atoms, where the nitrogen and sulfur heteroatoms may also optionally be oxidized. The heterocyclic group may be attached at a heteroatom or carbon atom.

Exemplary monocyclic heterocyclic groups include pyrrolidinyl, pyrrolyl, pyrazolyl, oxetanyl, pyrazolinyl, imidazolyl, tetrazolyl, imidazolinyl, imidazolidinyl, oxazolyl, oxazolidinyl, isoxazolinyl, isoxazolyl, thiazolyl, thiadiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, furyl, tetrahydrofuryl, thienyl, oxadiazolyl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, 4-piperidonyl, pyridyl, 2-pyridone, N-lower alkyl-pyridone, e.g. N-lower alkyl-2-pyridone, pyrazinyl, pyrimidinyl, pyridazinyl, tetrahydropyranlyl, morpholinyl, thiamorpholinyl, S-oxo-thiamorpholinyl, S,S-dioxo-thiamorpholinyl, 1,3-dioxolane and tetrahydro-1,1-dioxothienyl, and the like.

Exemplary bicyclic heterocyclic groups include indolyl, benzothiazolyl, benzoxazolyl, benzothienyl, quinuclidinyl, quinolinyl, tetrahydroisoquinolinyl, isoquinolinyl, benzimidazolyl, benzopyranlyl, indolizinyl, benzofuryl, chromonyl, coumarinyl, benzopyranlyl, cinnolinyl, quinoxalinyl, indazolyl, pyrrolopyridyl, furopyridinyl (such as furo[2,3-c]pyridinyl, furo[3,2-b]-

0956960-092801

pyridinyl] or furo[2,3-b]pyridinyl), dihydroisoindolyl, dihydroquinazolinyl (such as 3,4-dihydro-4-oxo-quinazolinyl) and the like.

Exemplary tricyclic heterocyclic groups include carbazolyl, benzindolyl, phenanthrolinyl, acridinyl, phenanthridinyl, xanthenyl and the like.

The term "heterocyclyl" includes substituted heterocyclic groups. Substituted heterocyclic groups refer to heterocyclic groups substituted with 1, 2 or 3 of the following:

- (a) alkyl;
- (b) hydroxy (or protected hydroxy);
- (c) halo;
- (d) oxo (i.e. = O);
- (e) amino, alkylamino or dialkylamino;
- (f) alkoxy;
- (g) cycloalkyl;
- (h) carboxy;
- (i) heterocycloxy;
- (j) alkoxycarbonyl, such as unsubstituted lower alkoxycarbonyl;
- (k) mercapto;
- (l) nitro;
- (m) cyano;
- (n) sulfonamido, sulfonamidoalkyl or sulfonamidodialkyl;
- (o) aryl;
- (p) alkylcarbonyloxy;
- (q) arylcarbonyloxy;
- (r) arylthio;
- (s) aryloxy;
- (t) alkylthio;
- (u) formyl;
- (v) aralkyl; or

0996990099690

The term "heterocyclooxy" denotes a heterocyclic group bonded through an oxygen bridge.

The term "heteraryloxy" refers to heteroaryl-O-.

The term "heteroaroyl" refers to heteroaryl-C(O)-.

Encompassed by the invention are prodrug derivatives, e.g., any pharmaceutically acceptable prodrug ester derivatives of the carboxylic acids of the invention (-COR1 being carboxy) which are convertible by solvolysis or under physiological conditions to the free carboxylic acids.

Examples of such carboxylic acid esters include esters defined by -C(O)R₁, and are preferably lower alkyl esters, cycloalkyl esters, lower alkenyl esters, benzyl esters, mono or disubstituted lower alkyl esters, e.g. the ω-(amino, mono- or di-lower alkylamino, carboxy, lower alkoxy-carbonyl)-lower alkyl esters, the α-(lower alkanoyloxy, lower alkoxy-carbonyl or di-lower alkylaminocarbonyl)-lower alkyl esters, such as the pivaloyloxy-methyl ester, and the like conventionally used in the art.

Preferred meanings of R are hydrogen or lower alkyl;

Preferred meanings of Q are $-C(O)R_1$ wherein R_1 is hydroxy, lower alkoxy or aryloxy.

Preferred meanings of R_2 are hydrogen, halogen or lower alkyl.

Preferred meanings of R_3 are halogen or lower alkyl.

Preferred meanings of R_4 are phenyl or phenyl substituted by one or more substituents selected from the group consisting of lower alkyl, lower alkoxy, halogen and trifluoromethyl.

Preferred meaning of R_5 is hydrogen.

Preferred meanings of R_6 are phenyl or phenyl substituted by one or more substituents selected from the group consisting of lower alkyl, lower alkoxy, halogen and trifluoromethyl.

Preferred W is O.

Preferred X is $-S(O)_2R_4$ or $-S(O)_2NR_5R_6$

Preferred Y is O.

Preferred Z is hydroxy.

The integer "n" preferably is zero, 1 or 2.

The compounds of the invention depending on the nature of the substituents, may possess one or more asymmetric centers. The resulting diastereoisomers, enantiomers and geometric isomers are encompassed by the instant invention. The present invention further includes tautomers wherein groups possible for tautomerism are involved.

Preferred are the compounds of formula I in which

W is O or S;

X is $-S(O)_2R_4$; R_4 being lower alkyl, phenyl or phenyl substituted by one or more substituents selected from the group consisting of lower alkyl, lower alkoxy, halogen and trifluoromethyl; or is $-S(O)_2NR_5R_6$ or is $-C(O)NR_5R_6$; R_5 , in each case, being hydrogen or lower alkyl and R_6 , in each case, being hydrogen, lower alkyl, lower alkyl substituted by NR_5R_6 , 3- to 7-membered cycloalkyl, phenyl, phenyl substituted by one or more substituents selected from the group consisting of lower alkyl, lower alkoxy, halogen and trifluoromethyl; pyridyl or N-lower alkyl-2-pyridone; or R_5 and R_6 combined, in each case, being alkylene or alkylene interrupted by O or $S(O)_2$ which together with the nitrogen atom to which they are attached form

0996960-092804

a 5- to 7- membered ring;

Y is O or H₂;

Z is hydrogen, halogen, hydroxy, alkoxy, aralkoxy, acyloxy or alkoxy-carbonyloxy;

R is hydrogen;

Q is -C(O)R₁ wherein R₁ is hydroxy, lower alkoxy or -NR₅R₆; R₅ being hydrogen or lower alkyl and R₆ being hydrogen, lower alkyl, lower alkoxy or R₅ and R₆ combined being alkylene or alkylene interrupted by O which together with the nitrogen atom to which they are attached form a 5- to 7- membered ring ;


R₂ is hydrogen, halogen or lower alkyl;

R₃ is halogen or lower alkyl;

n represents zero, 1 or 2;

and pharmaceutically acceptable salts thereof.

Preferred are the compounds of the formula



in which

W is O or S;

X is -SR₄, -S(O)R₄, -S(O)₂R₄, -S(O)₂NR₅R₆ or -C(O)NR₅R₆;

Y is O or H₂;

Z is hydrogen, halogen, hydroxy, alkoxy, aralkoxy, acyloxy or alkoxy-carbonyloxy;

R₁ is hydroxy, lower alkoxy or aryloxy;

R₂ is hydrogen, halogen or lower alkyl;

R₃ is halogen or lower alkyl;

R₄ is optionally substituted alkyl, aryl, aralkyl, heteroaryl or heteroaralkyl;


R5, R6 and R7 are independently hydrogen, optionally substituted alkyl, cycloalkyl, aryl, aralkyl, heteroaryl, or heteroaralkyl; or R5 and R6 combined are alkylene optionally interrupted by O, S, S(O), S(O)₂ or NR7 which together with the nitrogen atom to which they are attached form a 5- to 7- membered ring;

n represents zero, 1 or 2;

and pharmaceutically acceptable salts thereof.

Preferred are compounds of formula IA, and pharmaceutically acceptable salts thereof, wherein Z is different from hydrogen when X is -C(O)NR5R6.

Preferred are also the compounds of the formula



in which

W is O or S;

X is -SR4, -S(O)R4, -S(O)₂R4, -S(O)₂NR5R6 or -C(O)NR5R6;

Y is O or H₂;

Z is hydrogen, halogen, hydroxy, alkoxy, aralkoxy, acyloxy or alkoxycarbonyloxy;

Q is 5-tetrazolyl;

R2 is hydrogen, halogen or lower alkyl;

R3 is halogen or lower alkyl;

R4 is optionally substituted alkyl, aryl, aralkyl, heteroaryl or heteroaralkyl;

R5, R6 and R7 are independently hydrogen, optionally substituted alkyl, cycloalkyl, aryl, aralkyl, heteroaryl, or heteroaralkyl; or R5 and R6 combined are alkylene optionally interrupted by O, S, S(O), S(O)₂ or NR7 which together with the nitrogen atom to which they are attached


09699660-09699660

form a 5- to 7- membered ring;

n represents zero, 1 or 2;

and pharmaceutically acceptable salts thereof.

Further preferred are the compounds of the formula



in which

X is $-S(O)_2R_4$, $-S(O)_2NR_5R_6$ or $-C(O)NR_5R_6$;

Z is hydroxy, lower alkanoyloxy or lower alkoxy;


R1 is hydroxy or lower alkoxy;

R2 and R3 are lower alkyl;

R4 is aryl;

R5, R6 and R7 are independently hydrogen, optionally substituted alkyl, cycloalkyl, aryl, aralkyl, heteroaryl, or heteroaralkyl; or R5 and R6 combined are alkylene optionally interrupted by O, S, $S(O)$, $S(O)_2$ or NR_7 which together with the nitrogen atom to which they are attached form a 5- to 7- membered ring; and pharmaceutically acceptable salts thereof.

Further preferred are also the compounds of the formula



in which

X is $-S(O)_2R_4$, $-S(O)_2NR_5R_6$ or $-C(O)NR_5R_6$;

Z is hydroxy, lower alkanoyloxy or lower alkoxy;

Q is 5-tetrazolyl;


R2 and R3 are lower alkyl;

R4 is aryl;

R5, R6 and R7 are independently hydrogen, optionally substituted alkyl, cycloalkyl, aryl, aralkyl, heteroaryl, or heteroaralkyl; or R5 and R6 combined are alkylene optionally interrupted by O, S, S(O), S(O)₂ or NR7 which together with the nitrogen atom to which they are attached form a 5- to 7- membered ring; and pharmaceutically acceptable salts thereof.

Preferred are compounds of formula I, IA, IB and IB', and pharmaceutically acceptable salts thereof, wherein X is -S(O)₂R4 or -S(O)₂NR5R6.

Most preferred are the compounds of the formula




in which

X is -S(O)₂R4 or -S(O)₂NR5R6;

R4 is monocyclic aryl;

R5, R6 and R7 are independently hydrogen, optionally substituted alkyl or aryl; or R5 and R6 combined are CH₂CH₂-U-CH₂CH₂ wherein U is CH₂, O, NR7, S, S(O) or S(O)₂ which together with the nitrogen atom to which they are attached form a 6-membered ring; pharmaceutically acceptable prodrug esters thereof; and pharmaceutically acceptable salts thereof.

Most preferred are also the compounds of the formula



in which

X is $-\text{S}(\text{O})_2\text{R}_4$ or $-\text{S}(\text{O})_2\text{NR}_5\text{R}_6$;

Q is 5-tetrazolyl;

R₄ is monocyclic aryl;

R₅, R₆ and R₇ are independently hydrogen, optionally substituted alkyl or aryl; or R₅ and R₆ combined are $\text{CH}_2\text{CH}_2\text{-U-CH}_2\text{CH}_2$ wherein U is CH_2 , O, NR₇, S, S(O) or S(O)₂ which together with the nitrogen atom to which they are attached from a 6-membered ring; pharmaceutically acceptable prodrug esters thereof; and pharmaceutically acceptable salts thereof.


Particularly preferred are the compounds of formula IC and IC' wherein X is S(O)₂R₄ and R₄ is phenyl optionally substituted by lower alkyl, halo, lower alkoxy or trifluoromethyl; pharmaceutically acceptable salts thereof; and prodrug derivatives thereof.

Pharmaceutically acceptable salts of any acidic compounds of the invention are salts formed with bases, namely cationic salts such as alkali and alkaline earth metal salts, such as sodium, lithium, potassium, calcium, magnesium, as well as ammonium salts, such as ammonium, trimethylammonium, diethylammonium, and tris-(hydroxymethyl)-methylammonium salts.


Similarly acid addition salts, such as of mineral acids, organic carboxylic, and organic sulfonic acids e.g. hydrochloric acid, methanesulfonic acid, maleic acid, are possible provided a basic group, such as pyridyl, constitutes part of the structure.

096600-096600

Compounds of formula I may be prepared from appropriately substituted phenols of the formula




wherein R2 and R3 have meaning as defined herein and W is oxygen (prepared according to methods well-known in the literature) by first converting the hydroxyl group to a leaving group L, such as trifluoromethylsulfonyloxy or halo, such as chloro or fluoro, to form compounds of the formula




wherein R2 and R3 have meanings as defined herein.

Compounds of formula III wherein L is trifluoromethylsulfonyloxy can be converted to compounds of formula III wherein L is e.g. chloro by treating the former with lithium chloride in an inert solvent such as N-methylpyrrolidone, N,N-dimethylformamide or dimethylsulfoxide at a temperature ranging from room temperature to about 130°C.

Compounds of formula III can be converted to compounds of the formula




by reaction with appropriately substituted phenols or thiophenols of the formula



wherein R has meaning as defined herein, X' and Z' represent X and Z as defined herein, or X' and Z' are groups convertible to X and Z respectively, in the presence of a base such as sodium hydride or potassium carbonate in an inert solvent such as N-methylpyrrolidone, N,N-dimethylformamide or dimethylsulfoxide at a temperature ranging from room temperature to about 130°C. Compounds of formula V can be obtained using methodologies described herein or in the art.


Alternatively, compounds of formula IV can be obtained by condensing bisaryl iodonium tetrafluoroborates of the formula



in which R, X' and Z' are as defined above, with phenols or thiophenols of formula II wherein W is oxygen or sulfur as described in the art, e.g. in the presence of a copper catalyst and a base such as triethylamine in an inert solvent such as dichloromethane.


Compounds of formula IV wherein Z' is alkoxy or aralkoxy can be converted to compounds of formula IV wherein Z' is hydroxy according to methods well-known in the art, e.g. using acid such as hydrobromic acid or a boron trihalide, such as boron trichloride or boron tribromide when Z' is in particular a methoxy group or using hydrogen in the presence of a catalyst such as palladium on carbon when Z' is in particular a benzyloxy group.

Compounds of the invention wherein X is -S(O)₂NR₅R₆ may be prepared by, for example, first treating compounds of formula IV, wherein R and X' are hydrogen and X' is located at the 3' position, and Z' is hydroxy, alkoxy or aralkoxy and Z' is located at the 4' position, with chlorosulfonic acid in an organic solvent such as dichloromethane to produce compounds of the formula




wherein Z' is as defined above. Compounds of formula VII in which Z' is hydroxy, may be converted to compounds of formula VII wherein Z' is a protected hydroxyl group such as alkoxy, aralkoxy, alkanoyloxy, alkoxycarbonyloxy or trialkylsiloxy using methods and conditions well-known in the art.

Compounds of formula VII wherein Z' is alkoxy, aralkoxy, alkanoyloxy, alkoxycarbonyloxy or trialkylsiloxy can be converted to compounds of the formula




wherein Z' is as defined above by reaction with a chlorinating agent such as oxalyl chloride or thionyl chloride in an inert solvent such as dichloromethane or tetrahydrofuran in the presence of a catalytic amount of N,N-dimethylformamide.

A reaction of compounds of formula VIII with primary or secondary amines of formula R₅R₆NH wherein R₅ and R₆ are as defined herein in the presence of a base such as N-methylmorpholine or triethylamine in an organic solvent such as dichloromethane affords compounds of the formula




Compounds of formula IX wherein Z' is as defined above can be converted to compounds of formula IX wherein Z' is hydroxy using methods and conditions well-known in the art or as illustrated herein.

Conversion of nitro substituted compounds of formula IV, for example those of formula IX, to amines of e.g. the formula




can be achieved according to methods described in the art, e.g. with hydrogen in the presence of a catalyst such as palladium on carbon in a polar solvent such as ethanol or tetrahydrofuran.

Any resulting amines, for example those of formula X, can be treated with acylating agents such as ethyl oxalyl chloride, ethyl malonyl chloride, ethyl succinyl chloride or ethyl bromoacetate in the presence of a base such as N-methylmorpholine or triethylamine in an organic solvent such as dichloromethane, tetrahydrofuran or N,N-dimethylformamide to form compounds of formula I, e.g. those of the formula



wherein R1 is alkoxy, Y is oxygen or H₂, Z' represents Z as defined herein or Z' is a group convertible to Z and n represents an integer from zero to four. Compounds of formula I wherein Q is -C(O)R1 and R1 is alkoxy can also be prepared by condensing e.g. compounds of formula X with acylating agents such as dimethyl or diethyl oxalate at elevated temperature using the acylating agent both as a reagent and a solvent.


Compounds of formula I wherein Q is 5-tetrazolyl, Y is oxygen and n is zero can be obtained by condensing e.g. amines of formula X with 1H-tetrazole-5-carbonyl chloride (prepared from 1H-tetrazole-5-carboxylic acid ethyl ester using methods known in the art) in the presence of a base such as pyridine or triethylamine in an organic solvent such as acetonitrile or N,N-dimethylformamide to give compounds of formula I, e.g. those of formula



Compounds of formula I wherein Q is 5-tetrazolyl, Y is oxygen or H₂ and n is an integer from 1 to 4 may be prepared according to methods well-known in the art.


Compounds of formula I wherein Q is -C(O)R₁ and R₁ is e.g. alkoxy or aryloxy can be hydrolyzed to compounds of formula I wherein R₁ is hydroxy according to conventional methods, e.g. using an aqueous base such as an alkali metal carbonate or hydroxide in an organic solvent such as ethanol or tetrahydrofuran.

Similarly, other compounds of formula IV wherein, X' and Z' represent X and Z as defined herein or X' and Z' are groups convertible to X and Z respectively are converted to compounds of the formula




in a manner illustrated herein or by modifications thereof and, if necessary, said compounds are converted to the corresponding compounds of formula I, by converting X' and Z' to X and Z respectively. For example, compounds wherein X' is -COOH are converted to the corresponding amides of formula I wherein X is -C(O)NR₅R₆ according to methods well known in the art. Analogously, compounds of formula I wherein Q is -C(O)R₁ and R₁ is lower alkoxy or hydroxy can be converted to compounds wherein -C(O)R₁ is -C(O)NR₅R₆.

Compounds of the invention wherein X is -S(O)₂R₄, for example compounds of the formula




in which R4 is optionally substituted alkyl, aryl, aralkyl, heteroaryl or heteroaralkyl; R1 is hydroxy, optionally substituted alkoxy, aryloxy, heteroaryloxy, aralkoxy, cycloalkoxy or heteroaralkoxy; R2 is hydrogen, halogen or lower alkyl; R3 is halogen or lower alkyl; and n represents zero or an integer from 1 to 4; or a pharmaceutically acceptable salt thereof; can be prepared using the following sequence of steps:

(a) condensing 1,4-benzoquinone in an organic solvent such as dichloromethane, preferably at room temperature, with a sulfinic acid of the formula $R_4S(O)OH$ to obtain a compound of the formula




in which R4 has meaning as defined herein, and the sulfinic acid may be prepared by reduction of a sulfonyl chloride of formula $R_4S(O)_2Cl$ using a reducing agent such as sodium borohydride in an inert solvent such as tetrahydrofuran, or alternatively using sodium bisulfite as the reducing agent in the presence of an aqueous base such as sodium hydroxide;


(b) reacting a compound of formula XII in the presence of a base such as sodium hydride or potassium carbonate in an inert solvent such as N-methylpyrrolidone, N,N-dimethylformamide or dimethylsulfoxide at a temperature ranging from room temperature to about 130°C with a compound of the formula



in which R2, R3 and L have meaning as defined herein to obtain a compound of the formula



(c) converting the nitro compound of formula IXA according to methods well described in the art, e.g. with hydrogen in the presence of a catalyst such as palladium on carbon in a polar solvent such as ethanol or tetrahydrofuran, to a corresponding amine of the formula



in which R2, R3 and R4 have meaning as defined above;

(d) condensing the amine of formula XA in the presence of a base such as N-methylmorpholine or triethylamine in an organic solvent such as dichloromethane, tetrahydrofuran or N,N-dimethylformamide with a reactive functional derivative of an acid corresponding to the formula




In which R1 is optionally substituted alkoxy, aryloxy, heteroaryloxy, aralkoxy, cycloalkoxy or heteroaralkoxy; and n has meaning as defined above, e.g. an acid halide such as ethyl oxalyl chloride, ethyl malonyl chloride, ethyl succinyl chloride, diethyl and dimethyl oxalate, to obtain an ester of formula XIB wherein R1 and n are as defined for formula XIII; and R2, R3 and R4 have meaning as defined above; and if desired

(e) hydrolyzing said ester to a carboxylic acid of formula XIB in which R1 is hydroxy according to conventional methods, e.g. using an aqueous base such as an alkali metal carbonate or hydroxide in an organic solvent such as ethanol or tetrahydrofuran; and if desired converting said acid to a pharmaceutically acceptable salt thereof.

103250-03599600

Compounds of formula I wherein Q is 5-tetrazolyl, Y is oxygen and n is zero can be obtained by condensing e.g. amines of formula XA with 1H-tetrazole-5-carbonyl chloride in the presence of a base such as pyridine or triethylamine in an organic solvent such as acetonitrile or N,N-dimethylformamide to give compounds of formula I, e.g. those of formula



wherein R2, R3 and R4 are as defined above.

The starting compounds and intermediates which are converted to the compounds of the invention in a manner described herein, functional groups present, such as amino, thiol, carboxyl, and hydroxy groups, are optionally protected by conventional protecting groups that are common in preparative organic chemistry. Protected amino, thiol, carboxyl, and hydroxy groups are those that can be converted under mild conditions into free amino, thiol, carboxyl and hydroxy groups without the molecular framework being destroyed or other undesired side reactions taking place.

The purpose of introducing protecting groups is to protect the functional groups from undesired reactions with reaction components under the conditions used for carrying out a desired chemical transformation. The need and choice of protecting groups for a particular reaction is known to those skilled in the art and depends on the nature of the functional group to be protected (hydroxy group, amino group, etc.), the structure and stability of the molecule of which the substituent is a part and the reaction conditions.

Well-known protecting groups that meet these conditions and their introduction and removal are described, for example, in J.F.W. McOmie, "Protective Groups in Organic Chemistry", Plenum Press, London, New York, 1973, T. W. Greene, "Protective Groups in Organic Synthesis", Wiley, New York, 1991.

The above-mentioned reactions are carried out according to standard methods, in the presence or absence of diluent, preferably such as are inert to the reagents and are solvents thereof, of catalysts, condensing or said other agents respectively and/or inert atmospheres, at low temperatures, room temperature or elevated temperatures (preferably at or near the boiling point of the solvents used), and at atmospheric or super-atmospheric pressure. The preferred solvents, catalysts and reaction conditions are set forth in the appended illustrative Examples.

The invention further includes any variant of the present processes, in which an intermediate product obtainable at any stage thereof is used as starting material and the remaining steps are carried out, or in which the starting materials are formed in situ under the reaction conditions, or in which the reaction components are used in the form of their salts or optically pure antipodes.

Compounds of the invention and intermediates can also be converted into each other according to methods generally known per se.

The invention also relates to any novel starting materials and processes for their manufacture.

Depending on the choice of starting materials and methods, the new compounds may be in the form of one of the possible isomers or mixtures thereof, for example, as substantially pure geometric (cis or trans) isomers, optical isomers (antipodes), racemates, or mixtures thereof. The aforesaid possible isomers or mixtures thereof are within the purview of this invention.

Any resulting mixtures of isomers can be separated on the basis of the physico-chemical differences of the constituents, into the pure geometric or optical isomers, diastereoisomers, racemates, for example by chromatography and/or fractional crystallization.

Any resulting racemates of final products or intermediates can be resolved into the optical antipodes by known methods, e.g. by separation of the diastereoisomeric salts thereof, obtained with an optically active acid or base, and liberating the optically active acidic or basic compound. The carboxylic acid intermediates can thus be resolved into their optical antipodes e.g. by fractional crystallization of d- or l-(alpha-methylbenzylamine, cinchonidine, cinchonine, quinine, quinidine, ephedrine, dehydroabietylamine, brucine or strychnine)-salts. Racemic products can also be resolved by chiral chromatography, e.g. high pressure liquid chromatography using a chiral adsorbent.

Finally, compounds of the invention are either obtained in the free form, or as a salt thereof if salt forming groups are present.

Acidic compounds of the invention may be converted into salts with pharmaceutically acceptable bases, e.g. an aqueous alkali metal hydroxide, advantageously in the presence of an ethereal or alcoholic solvent, such as a lower alkanol. From the solutions of the latter, the salts may be precipitated with ethers, e.g. diethyl ether. Resulting salts may be converted into the free compounds by treatment with acids. These or other salts can also be used for purification of the compounds obtained.

Compounds of the invention having basic groups can be converted into acid addition salts, especially pharmaceutically acceptable salts. These are formed, for example, with inorganic acids, such as mineral acids, for example sulfuric acid, a phosphoric or hydrohalic acid, or with organic carboxylic acids, such as (C₁-C₄)-alkanecarboxylic acids which, for example, are unsubstituted or substituted by halogen, for example acetic acid, such as saturated or unsaturated dicarboxylic acids, for example oxalic, succinic, maleic or fumaric acid, such as hydroxy-carboxylic acids, for example glycolic, lactic, malic, tartaric or citric acid, such as amino acids, for example aspartic or glutamic acid, or with organic sulfonic acids, such as (C₁-C₄)-alkyl-sulfonic acids (for example methanesulfonic acid) or arylsulfonic acids which are

T03260-0969660

In view of the close relationship between the free compounds and the compounds in the form of their salts, whenever a compound is referred to in this context, a corresponding salt is also intended, provided such is possible or appropriate under the circumstances.

The pharmaceutical compositions according to the invention are those suitable for enteral, such as oral or rectal, transdermal and parenteral administration to mammals, including man, for the treatment of diseases associated with an imbalance of thyroid hormones, such as hypo- and hyperthyroidism, obesity, osteoporosis, depression, and in particular for the treatment and/or pre-vention of occlusive cardiovascular conditions in which hyperlipidimia and hyperlipoproteinemia are implicated, comprising an effective amount of a pharmacologically active compound of the invention, alone or in combination with one or more pharmaceutically acceptable carriers.

The pharmacologically active compounds of the invention are useful in the manufacture of pharmaceutical compositions comprising an effective amount thereof in conjunction or admixture with excipients or carriers suitable for either enteral or parenteral application. Preferred are tablets and gelatin capsules comprising the active ingredient together with a) diluents, e.g. lactose, dextrose, sucrose, mannitol, sorbitol, cellulose and/or glycine; b) lubricants, e.g. silica, talcum, stearic acid, its magnesium or calcium salt and/or polyethyleneglycol; for tablets also c) binders e.g. magnesium aluminum silicate, starch paste, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose and or polyvinylpyrrolidone; if desired d) disintegrants, e.g. starches, agar, alginic acid or its sodium salt, or effervescent mixtures; and/or e) absorbants, colorants, flavors and sweeteners. Injectable compositions are preferably aqueous isotonic solutions or suspensions, and suppositories are advantageously prepared from fatty emulsions or suspensions. Said compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution

promoters, salts for regulating the osmotic pressure and/or buffers. In addition, they may also contain other therapeutically valuable substances. Said compositions are prepared according to conventional mixing, granulating or coating methods, respectively, and contain about 0.1 to 75%, preferably about 1 to 50%, of the active ingredient.

Suitable formulations for transdermal application include an effective amount of a compound of the invention with carrier. Advantageous carriers include absorbable pharmacologically acceptable solvents to assist passage through the skin of the host. Characteristically, transdermal devices are in the form of a bandage comprising a backing member, a reservoir containing the compound optionally with carriers, optionally a rate controlling barrier to deliver the compound of the skin of the host at a controlled and predetermined rate over a prolonged period of time, and means to secure the device to the skin.

A unit dosage for a mammal of about 50 to 70 kg may contain between about 0.01 mg and 10 mg of the active ingredient. The dosage of active compound is dependent on the species of warm-blooded animal (mammal), the body weight, age and individual condition, on the form of administration, and on the compound involved.

The problem solved by the present invention is to provide pharmacologically potent hypolipidemic agents which reduce plasma cholesterol levels in mammals. The compounds of the invention demonstrate potent binding to the triiodothyronine (T_3) nuclear receptor which is indicative of upregulation of LDL receptor activity and enhancement of the clearance of LDL-cholesterol from the circulation. The compounds of the invention are thus particularly useful in mammals as hypocholesteremic agents for the treatment and prevention of occlusive cardiovascular conditions in which hypercholesteremia is implicated, by reducing plasma levels of total and LDL-cholesterol. The invention furthermore relates to the use of the compounds according to the invention for the preparation of medicaments, in particular of medicaments useful for the treatment and prevention of occlusive cardiovascular conditions in which hypercholesteremia is implicated, by reducing plasma levels of total and LDL-cholesterol.

Compounds of the invention also reduce lipoprotein (a) levels and are thus useful for the treatment and prevention of occlusive cardiovascular conditions in which Lp(a) is implicated.

09066960-092301

Selective thyromimetic hypolipidemic agents of the invention which are substantially free of undesirable cardiac side effects associated with thyroid hormones are preferred.

The above-cited properties are demonstrable in vitro and in vivo tests, using advantageously mammals, e.g. mice, rats, dogs, monkeys or isolated organs, tissues and preparations thereof. Said compounds can be applied in vitro in the form of solutions, e.g. preferably aqueous solutions, and in vivo either enterally, parenterally, advantageously intravenously, e.g. as a suspension or in aqueous solution. The dosage in vitro may range between about 10^{-7} molar and 10^{-11} molar concentrations. The dosage in vivo may range depending on the route of administration, between about 0.1 and 1000 micrograms/kg, preferably between about 0.5 and 300 micrograms/kg, advantageously between about 1 and 100 micrograms/kg.

The compounds of the invention bind to the triiodothyronine (T_3) receptor and are thus useful as thyroid hormone agonists in mammals.

The in vitro binding to T_3 nuclear receptors is determined as follows:

Rat liver nuclei and plasma membrane preparations are obtained from Sprague-Dawley (CD) rats (Charles River Labs.) by differential centrifugation as described by Emmelot et al (Methods in Enzymology 31:75, Part A, 1974) with minor modifications. The nuclear fraction obtained from the 275 x g pellet is further purified as generally described by Spindler et al (J. Biol. Chem. 250:4118, 1975).

The novel test compounds are assayed for binding to the nuclei by the method of Spindler et al (J. Biol. Chem. 250:4118, 1975). The nuclei are incubated at 22°C with 0.3 nM of [^{125}I]-L-triiodothyronine (L-T_3). Parallel incubations are conducted with tubes containing, in addition to the nuclei and radioactive L-T_3 , either various concentrations of the test compounds or 3 μM of nonradioactive L-T_3 . The latter is used as a measure of nonspecific binding. The radio-activity bound to the nuclei is determined following centrifugation of the reaction mixture at 800 x g for 7 minutes and washing of the pellet obtained. The amount of [^{125}I]- L-T_3 specifically bound is determined by subtracting the amount nonspecifically bound (radioactivity contained in

the nuclear pellet following incubation with excess (3 μ M) non-radioactive L-T₃. The concentration of test compound which inhibits the specific binding of [¹²⁵I]-L-T₃ by 50 percent (IC₅₀) is determined graphically from the reciprocal plot of the specifically bound [¹²⁵I]-L-T₃ versus the various concentrations of the test compound.

Cholesterol lowering activity is determined in the rat as follows:

Male Sprague-Dawley rats (230-250 g) (Taconic Farms) are maintained ad libitum on water and a high cholesterol diet (1.5% cholesterol and 0.5% cholic acid) for two weeks prior to and during the 7-day treatment period. Groups of animals are treated orally by gavage with the vehicle alone or with test compound for 7 consecutive days. After the last dose, animals are fasted for 18 hours and blood is collected. Blood samples are centrifuged at 2500 rpm for 10 minutes to prepare plasma for total cholesterol determination as well as LDL and HDL cholesterol concentrations. HDL values are determined after LDL/VLDL precipitation (Warnick and Albers, 1978). All samples are analyzed enzymatically for cholesterol with a diagnostic reagent kit (Sigma Chemical Co., St. Louis, MO). The analysis is performed on a Bio-Mek automated work station. LDL/VLDL fractions are precipitated in the following manner: 0.35 mL of plasma is aliquoted into Eppendorf tubes to which 12 μ L of 2M manganese chloride, 11.2 μ L of sodium heparin (Porcine Intestinal, 5000 units/mL), and 8.3 μ L of normal saline are added. The samples are vortexed and are placed on ice for 15 minutes, then centrifuged at 4°C for 10 minutes at 1300 rpm and the supernatant is enzymatically analyzed for cholesterol. The HDL cholesterol concentration is adjusted for dilution by multiplying the supernatant cholesterol value by 1.09. LDL/VLDL cholesterol values are obtained by subtracting HDL cholesterol from total cholesterol.

Cholesterol lowering activity can also be evaluated in normocholesterolemic dogs fed regular chow following the procedure described above, by administration of test compound orally for 5 days.

Cholesterol and Lp(a) lowering activity can be determined in the normolipemic cynomolgus monkey as follows:

Adult male and female cynomolgus monkeys (*Macaca fascicularis*) weighing 3-7 kg are used. Animals are individually housed and fed a standard monkey chow diet (Purina 5047)


supplemented with fresh fruits and vegetables. Each animal serves as its own control and each dosing regime is followed by a wash out period. Test compounds were dissolved in ethanol, imbibed in fruit pulp and administered orally. Animals are dosed once a day with test compounds for a treatment period varying from 8 to 28 days. Blood samples are obtained after an overnight fast at baseline and at the end of the study. Blood samples (3 ml) are collected into Vacutainer tubes (containing EDTA) from the femoral vein of mechanically restrained non-sedated animals. Blood is centrifuged at 2000 rpm for 20 min at 4 °C. Plasma samples are divided into aliquots and stored at -70 °C until analyzed. Plasma concentrations of total cholesterol (TC) and triglycerides (TG) are determined by enzymatic methods using commercial kits (Sigma Diagnostics). High-density lipoprotein cholesterol (HDL-C) concentration is measured after precipitation of apoB-containing lipoproteins. Since the fasting plasma of cynomolgus monkeys on a chow diet contains negligible amounts of cholesterol in very low-density lipoprotein, the low-density lipoprotein cholesterol (LDL-C) concentration is calculated by subtracting HDL-C from TC. The assays are performed in 96-well microtiter plates, which are read in a microplate spectrophotometer (Dynatech MR 5000). Plasma concentrations of Lp(a) are determined by a commercial Lp(a) ELISA (PerImmune, Inc.) using the kit controls and reference Lp(a) standard. The Lp(a) ELISA employs a monoclonal antibody against apolipoprotein(a) for capture and a polyclonal antibody against apolipoprotein B for detection. The assay is specific for Lp(a) and does not measure free apo(a), apoB or plasminogen. Quantification of Lp(a) is not affected by apo(a) size. The Lp(a) plasma concentration is reported as milligrams of total Lp(a) mass. Samples from each study are assayed in a single run.

Illustrative of the invention, the compound of example 26 demonstrates an IC_{50} of about 0.17 nM, the compound of example 28 demonstrates an IC_{50} of about 0.13 nM, the compound of example 35 demonstrates an IC_{50} of about 1.00 nM and the compound of example 39 demonstrates an IC_{50} of about 0.04 nM in the T_3 nuclear receptor binding assay. Furthermore, said compound of example 26 significantly lowers serum cholesterol at a daily dose of about 20 micrograms (μ g)/kg p.o. in the rat and about 10 μ g/kg p.o. in the dog. Furthermore, Lp(a) levels in the normolipemic cynomolgus monkeys are lowered by about 40% after 4 week treatment with the said compound of example 26 at a daily oral dose of 75 μ g/kg.

The following Examples are intended to illustrate the invention and are not to be construed as being limitations thereon. Temperatures are given in degrees Centigrade. If not mentioned otherwise, all evaporations are performed under reduced pressure, preferably between about 15 and 100 mmHg (= 20-133 mbar). The structure of final products, intermediates and starting materials is confirmed by standard analytical methods, e.g. microanalysis and spectroscopic characteristics (e.g. MS, IR, NMR). Abbreviations used are those conventional in the art.

Example 1

N-{4-[3-(2,2-Dimethylpropylsulfamoyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}-oxamic acid



A. 3,5-Dimethyl-4-(4'-methoxyphenoxy)nitrobenzene

A suspension of sodium hydride (NaH; 60% dispersion in mineral oil; 64.11 g, 1.603 mol) in 350 mL of N-methylpyrrolidone (NMP) is cooled to 0°C and treated with a solution of 4-methoxyphenol (208.4 g, 1.679 mol) over 30 min. The mixture is warmed to room temperature (RT) and after 30 min, 4-chloro-3,5-dimethylnitrobenzene (283.2 g, 1.526 mol; prepared by the method described by Yokoyama et. al. in EP580550) is added in one portion and the reaction is heated at 120°C for 2 h. The reaction is cooled to room temperature (RT) and quenched with water (1500 mL). The suspension is cooled to 0°C, stirred for 30 min, then filtered and the filtercake is washed with water and dried in vacuo. The crude product, ethyl acetate (EtOAc; 2100 mL) and charcoal (42.6 g) were heated to reflux and the solids were removed by filtration through celite while hot. The filtrate is concentrated under reduced pressure to ca. 800 mL and the resulting suspension is cooled to 0°C and stirred for 30 min. The product is collected by vacuum filtration, washed with cold EtOAc and dried in vacuo to afford 3,5-dimethyl-4-(4'-methoxyphenoxy)nitrobenzene: NMR (CDCl₃) 2.22 (s, 6H), 3.78 (s, 3H), 6.68 (d, 2H, J = 8.7), 6.82 (d, 2H, J = 8.7), 8.02 (s, 2H).

B. 3,5-Dimethyl-4-(4'-hydroxyphenoxy)nitrobenzene

A mixture of the title A compound, 3,5-dimethyl-4-(4'-methoxyphenoxy)nitrobenzene (16.4 g, 60 mmol), acetic acid (AcOH; 100 mL) and aqueous 48% hydrobromic acid (HBr; 100 mL) is heated at 120°C for 16 h. The mixture is cooled to RT, diluted with water (200 mL) and the precipitated product is collected by vacuum filtration, washed with water and hexanes and dried in vacuo to afford 3,5-dimethyl-4-(4'-hydroxyphenoxy)nitrobenzene: NMR (CDCl₃) 2.22 (s, 6H), 6.62 (d, 2H, J = 8.7), 6.77 (d, 2H, J = 8.7), 8.0 (s, 2H).

C. 5-(2,6-Dimethyl-4-nitrophenoxy)-2-hydroxybenzenesulfonic acid

A solution of the title B compound, 3,5-dimethyl-4-(4'-hydroxyphenoxy)nitrobenzene (7.86 g, 30.35 mmol) in 150 mL of dichloromethane (CH₂Cl₂) is treated with chlorosulfonic acid (2.4 mL, 36.42 mmol) at RT. After 16 h, the reaction mixture is concentrated and the residue is dissolved in small amount of CH₂Cl₂ (ca. 5 mL). The product is precipitated by addition of brine (100mL), collected by vacuum filtration, washed with water, hexanes and diethyl ether (Et₂O) and dried in vacuo to give 5-(2,6-dimethyl-4-nitrophenoxy)-2-hydroxy-benzenesulfonic acid: NMR (DMSO-d₆) 2.18 (s, 6H), 6.67-6.83 (m, 3H), 8.1 (s, 2H), 10.07 (s, 1H).

D. 2-Benzyloxy-5-(2,6-dimethyl-4-nitrophenoxy)benzenesulfonic acid, cesium salt

A solution of the title C compound, 5-(2,6-dimethyl-4-nitrophenoxy)-2-hydroxy-benzenesulfonic acid (6.78 g, 20 mmol) in 100 mL of tetrahydrofuran (THF) and 50 mL of N,N-dimethylformamide (DMF) is treated with cesium carbonate (15.6 g, 48 mmol) and benzyl bromide (7.1 mL, 60 mmol) at RT, then heated at 75°C for 48 h. The reaction mixture is cooled to RT and quenched with aqueous 1N hydrochloric acid (HCl; 100 mL), and THF is evaporated. The precipitated product is collected by vacuum filtration, washed with water, Et₂O, EtOAc and CH₂Cl₂ and dried to afford 2-benzyloxy-5-(2,6-dimethyl-4-nitrophenoxy)benzenesulfonic acid, cesium salt: NMR (DMSO-d₆) 2.18 (s, 6H), 5.10 (s, 2H), 6.74 (dd, 1H, J = 8.7, 3.8), 6.97 (d, 1H, J = 8.7), 7.11 (d, 1H, J = 3.8), 7.23-7.39 (m, 3H), 7.59 (d, 2H, J = 7.5), 8.12 (s, 2H).

E. 2-Benzyloxy-5-(2,6-dimethyl-4-nitrophenoxy)benzenesulfonyl chlorid

A suspension of the title D compound, 2-benzyloxy-5-(2,6-dimethyl-4-nitrophenoxy)-

0966960-09304

F. 2-Benzyloxy-5-(2,6-dimethyl-4-nitrophenoxy)-N-(2,2-dimethylpropyl)benzenesulfonamide


G. 5-(4-Amino-2,6-dimethylphenoxy)-2-hydroxy-N-(2,2-dimethylpropyl)benzenesulfonamide

A mixture of the title F compound, 2-benzyloxy-5-(2,6-dimethyl-4-nitrophenoxy)-N-(2,2-dimethylpropyl)benzenesulfonamide (1.22 g, 2.45 mmol) and palladium on activated carbon (10 wt. %; 250 mg) in 40 mL of THF is stirred under hydrogen atmosphere (H₂, 1 atm) for 8 h. The catalyst is removed by vacuum filtration through celite, washed with THF and the combined filtrate and washings are concentrated. The residue is suspended in CH₂Cl₂ and the product is collected by vacuum filtration, washed with CH₂Cl₂ and dried to give 5-(4-amino-2,6-dimethylphenoxy)-2-hydroxy-N-(2,2-dimethylpropyl)benzenesulfonamide: NMR (DMSO-d₆) 0.80 (s, 9H), 1.90 (s, 6H), 2.55 (d, 2H, J = 7.5), 4.90 (br s, 2H), 6.31 (s, 2H), 6.87 (br s, 1H), 6.93 (br s, 2H), 6.97 (t, 1H, J = 7.5), 10.13 (s, 1H).

H. N-{4-[3-(2,2-Dimethylpropylsulfamoyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}-oxamic acid

A solution of the title G compound, 5-(4-amino-2,6-dimethylphenoxy)-2-hydroxy-N-(2,2-dimethylpropyl)benzenesulfonamide (750 mg, 1.98 mmol) in 10 mL of THF is cooled to 0°C and treated sequentially with NMM (545 mL, 4.96 mmol) and ethyl oxalyl chloride (442 mL, 3.96 mmol). The reaction is warmed to RT and after 1h, quenched with water. The mixture is partitioned between water and EtOAc and the organic solution is washed with aqueous 1N HCl and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue is dissolved in 10 mL of THF, then treated with aqueous 1N lithium hydroxide (LiOH; 1.8 mL, 1.8 mmol) at RT. After 1h, the reaction is quenched with aqueous 1N HCl and the product is taken up in EtOAc, washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The product is triturated with hexanes, then with Et₂O and dried in vacuo to afford of N-{4-[3-(2,2-dimethylpropylsulfamoyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}oxamic acid: NMR(CDCl₃) 0.89 (s, 9H), 2.13 (s, 6H), 2.70 (d, 2H, J = 7.5), 5.22 (t, 1H, J = 7.5), 6.84 (d, 1H, J = 3.7), 7.04 (d, 1H, J = 8.7), 7.15 (dd, 1H, J = 8.7, 3.7), 7.33 (s, 2H), 8.54 (br s, 1H), 9.18 (s, 1H); IR (KBr) 1759, 1693; ESI-MS 449 [M-1].


The following additional compounds may be similarly prepared.

			
Example no.	NMR	IR (KBr)	ESI-MS
R5, R6			
2 R5 = H, R6 = Ph-	(DMSO-d ₆): 1.91 (s, 6H), 6.79-6.99 (m, 6H), 7.11-7.17 (m, 2H), 7.55 (s, 2H), 9.99 (s, 1H), 10.46 (s, 1H), 10.52 (s, 1H)	1207, 1481, 1689	455 [M-1]

3 R5 = H, R6 = p-FC ₆ H ₄ -	(DMSO-d ₆): 1.91 (s, 6H), 6.72 (d, 1H, J = 3.0), 6.89-7.03 (m, 6H), 7.55 (s, 2H), 9.96 (s, 1H), 10.54 (s, 1H), 10.66 (s, 1H)	1207, 1481, 1508, 1693	473 [M-1] ⁻
4 R5 = H, R6 = o-FC ₆ H ₄ -	(MeOH-d ₄): 1.98 (s, 6H), 6.81 (d, 1H, J = 2.6), 6.87-7.14 (m, 5H), 7.36 (td, 1H, J = 7.9, 1.9), 7.46 (s, 2H)	1209, 1481, 1500, 1689	473 [M-1] ⁻
5 R5 = H, R6 = m-FC ₆ H ₄ -	(DMSO-d ₆): 1.97 (s, 6H), 6.75-6.84 (m, 4H), 6.92 (d, 1H, J = 8.8), 7.04 (dd, 1H, J = 8.8, 3.1), 7.20 (app q, 1H, J = 7.0), 7.58 (s, 2H), 10.35 (s, 1H), 10.59 (s, 1H), 10.66 (s, 1H)	1238, 1475, 1706	473 [M-1] ⁻
6 R = H, R6 = p-MeOC ₆ H ₄ -	(DMSO-d ₆): 1.90 (s, 6H), 3.67 (s, 3H), 6.68 (d, 1H, J = 3), 6.73 (d, 2H, J = 9.00), 6.89-6.99 (m, 4H), 7.54 (s, 2H), 9.58 (s, 1H), 10.45 (s, 1H), 10.64 (s, 1H)	1209, 1475, 1514, 1688	485 [M-1] ⁻
7 R5 = H, R6 = p-FC ₆ H ₄ CH ₂ -	(DMSO-d ₆): 2.02 (s, 6H), 4.03 (d, 2H, J = 6.4), 6.80-6.89 (m, 3H), 7.01 (t, 2H, J = 4.0), 7.22 (dd, 2H, J = 6.3, 8.7), 7.56 (s, 2H), 7.77 (t, 1H, J = 6.4), 10.32 (s, 1H), 10.64 (s, 1H)	1224, 1481, 1695	487 [M-1] ⁻
8 R5 = Me-, R6 = Ph-	(CDCl ₃): 1.96 (s, 6H), 3.14 (s, 3H), 6.54 (d, 1H, J = 2.4), 6.84-6.85 (m, 2H), 6.88-7.03 (m, 2H), 7.20-7.28 (m, 5H)	1209, 1230, 1481, 1691	469 [M-1] ⁻
9 R5 = H, R6 = n-Pr-	(CDCl ₃): 0.87 (t, 3H, J = 7.5), 1.38-1.56 (m, 2H), 2.11 (s, 6H), 2.94 (app q, 2H, J = 7.5), 5.0 (t, 1H, J = 7.5), 6.84 (d, 1H, J = 3), 7.0 (d, 1H, J = 9), 7.08 (dd, 1H, J = 9, 3), 7.32 (s, 2H), 8.44 (br s, 1H), 9.0 (br s, 1H)	1214, 1485, 1693, 1751	421 [M-1] ⁻ 440 [M+NH ₄] ⁺
10 R5 = H, R6 = i-Pr-	(DMSO-d ₆): 0.95 (d, 6H, J = 7.5), 2.02 (s, 6H), 3.2-3.31 (m, 1H), 6.86 (br s, 1H), 6.97 (br s, 2H), 7.06 (d, 1H, J = 7.5), 7.57 (s, 2H), 10.32 (br s, 1H), 10.64 (br s, 1H)	1209, 1485, 1698, 1742	421 [M-1] ⁻
11 R5 = H, R6 = n-Bu-	(DMSO-d ₆): 0.72 (t, 3H, J = 7.5), 1.1-1.35 (m, 4H), 2.03 (s, 6H), 2.76 (app q, 2H, J = 7.5), 6.87 (br s, 1H), 6.97 (br s, 2H), 7.1 (t, 1H, J = 7.5), 7.58 (s, 2H)	1321, 1481, 1691	435 [M-1] ⁻ 454 [M+NH ₄] ⁺


T02260" 09090600

12 R5 = H, R6 = i-Bu-	(DMSO-d ₆): 0.75 (d, 6H, J = 8.2), 1.49-1.62 (m, 1H), 2.03 (s, 6H), 2.55 (app t, 2H, J = 6.8), 6.85 (br s, 1H), 6.96 (br s, 2H), 7.17 (t, 1H, J = 6.8), 7.57 (s, 2H), 10.25 (br s, 1H), 10.61 (br s, 1H)	1209, 1481, 1693, 1762	435 [M-1] ⁻
13 R5 = H, R6 = t-Bu-	(DMSO-d ₆): 1.02 (s, 9H), 2.03 (s, 6H), 6.81-7.02 (m, 4H), 7.56 (s, 2H), 10.20 (br s, 1H), 10.61 (br s, 1H)	1204, 1486, 1645, 1754	435 [M-1] ⁻
14 R5 = H, R6 = Cyclohexyl-	(CDCl ₃): 1.06-1.32 (m, 5H), 1.50-1.73 (m, 5H), 2.01 (s, 6H), 3.02-3.14 (m, 1H), 5.05 (d, 1H, J = 7.5), 6.82 (d, 1H, J = 3), 6.97 (d, 1H, J = 9), 7.10 (dd, 1H, J = 9, 3), 7.32 (s, 2H), 8.52 (br s, 1H), 9.06 (br s, 1H)	1209, 1481, 1695, 1741	461 [M-1] ⁻
15 R5 = Me-, R6 = Me-	(MeOH-d ₄): 2.07 (s, 6H), 2.64 (s, 6H), 6.86-6.94 (m, 3H), 7.39 (s, 2H)	1234, 1486, 1685	407 [M-1] ⁻
16 R5, R6 = -(CH ₂) ₄ -	(MeOH-d ₄): 1.76-1.80 (m, 4H), 2.08 (s, 6H), 3.21-3.29 (m, 4H), 6.88-6.95 (m, 3H), 7.40 (s, 2H)	1230, 1479, 1693	433 [M-1] ⁻
17 R5, R6 = -(CH ₂) ₅ -	(MeOH-d ₄): 1.49-1.58 (m, 6H), 2.13 (s, 6H), 3.08-3.10 (m, 4H), 6.92-7.02 (m, 3H), 7.53 (s, 2H)	1207, 1481, 1658	447 [M-1] ⁻
18 R5 = H, R6 = MeOCH ₂ CH ₂ -	(CDCl ₃): 2.06 (s, 6H), 3.03 (t, 2H, J = 5.5), 3.22 (s, 3H), 3.35 (t, 2H, J = 5.5), 6.85 (br s, 2H), 6.97 (br s, 1H), 7.34 (s, 2H)	1218, 1483, 1689	437 [M-1] ⁻
19 R5, R6 = -(CH ₂) ₂ O(CH ₂) ₂ -	(MeOH-d ₄): 2.13 (s, 6H), 3.12 (t, 4H, J=6), 3.68 (t, 4H, J = 6), 6.95-7.03 (m, 3H), 7.54 (s, 2H)	1479, 1711, 1745	449 [M-1] ⁻
20 R5, R6 = -(CH ₂) ₂ SO ₂ (CH ₂) ₂ -	(CDCl ₃): 2.12 (s, 6H), 3.10-3.13 (m, 4H), 3.74-3.76 (m, 4H), 6.92-6.99 (m, 3H), 7.41 (s, 2H)	1209, 1481, 1697	497 [M-1] ⁻

21 R5 = H, R6 = 3-Pyridyl	(DMSO-d ₆): 1.95 (s, 6H), 6.81 (d, 1H, J = 3), 6.92 (d, 1H, J = 9), 7.00 (dd, 1H, J = 9, 3), 7.26 (dd, 1H, J = 8, 5), 7.41-7.45 (m, 1H), 7.56 (s, 2H), 8.21 (dd, 1H, J = 5, 1.5), 8.27 (d, 1H, J = 2.5), 10.34 (s, 1H), 10.66 (s, 1H)	1481, 1500, 1689	473 [M-1] ⁻
22 R5 = H, R6 = 	(DMSO-d ₆): 1.88 (s, 6H), 3.75 (s, 3H), 6.64-6.69 (m, 2H), 6.93 (d, 1H, J = 9), 6.99 (dd, 1H, J = 9, 2.5), 7.32 (dd, 1H, J = 9, 2.5), 7.53 (s, 2H), 7.77 (d, 1H, J = 2.5), 9.74 (s, 1H), 10.56 (s, 1H), 10.61 (s, 1H)		488 [M+1] ⁺

Example 23


N-{4-[3-(4-Fluorophenylsulfamoyl)-4-hydroxyphenylsulfanyl]-3,5-dimethylphenyl}-oxamic acid



The title compound is prepared similarly to Example 1: NMR (DMSO-d₆) 2.21 (s, 6H), 6.87 (d, 1H, J = 8.3), 6.95-7.12 (m, 6H), 7.65 (s, 2H), 9.99 (s, 1H), 10.73 (s, 1H), 10.97 (s, 1H); IR (KBr) 1163, 1506, 1690; ESI-MS 489 [M-1]⁻, 508 [M+NH₄]⁺.

Example 24

N-{4-[3-(4-Fluorophenylsulfamoyl)phenoxy]-3,5-dimethylphenyl}oxamic acid




The title compound is similarly to Example 1: NMR (DMSO-d₆) 1.91 (s, 6H), 6.73-6.74 (m, 1H), 6.95-7.00 (m, 2H), 7.04-7.10 (m, 2H), 7.21 (dd, 1H, J = 8.2, 2.4), 7.35 (d, 1H, J = 8.0),

0996660-0996660

7.52 (app t, 1H, J = 8.0), 7.59 (s, 2H), 10.21 (s, 1H), 10.75 (s, 1H); IR (KBr) 1161, 1223, 1509, 1697; ESI-MS 457 [M-1]⁻, 476 [M+NH₄]⁺.

Example 25


N-{4-[3-(4-Fluorophenylsulfamoyl)-4-hydroxyphenoxy]-3-methylphenyl}oxamic acid



The title compound is prepared similarly to Example 1: NMR (DMSO-d₆) 2.06 (s, 3H), 6.65 (d, 1H, J = 8.8), 6.92-7.18 (m, 7H), 7.55 (m, 7H), 7.55 (dd, 1H, J = 8.8, 2.4), 7.70 (d, 1H, J = 2.4), 9.98 (s, 1H), 10.69 (s, 1H), 10.72 (s, 1H), 10.75 (s, 1H); IR (KBr) 1326, 1487, 1506, 1692; ESI-MS 457 [M-1]⁻.

Example 26

N-{4-[3-(4-Fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}oxamic acid



A. 4-Fluorobenzenesulfinic acid

A solution of 4-fluorobenzenesulfonyl chloride (2 g, 10.28 mmol) in 50 mL of THF (distilled from Na-benzophenone) is cooled to 0°C and sodium borohydride (1.9 g, 51.4 mmol) is added portionwise. The reaction is stirred at 0°C for 2 h, then warmed to RT, and after 2 h, quenched with water (5 mL). The solvent is evaporated and the aqueous residue is acidified by addition of aqueous 6N HCl. The product is taken up in EtOAc, washed with brine, dried over anhydrous Na₂SO₄ and concentrated to give 4-fluorobenzenesulfinic acid: NMR (DMSO-d₆) 7.12 (app t, 2H, J = 8.3), 7.5 (dd, 2H, J = 8.3, 6); ESI-MS 159 [M-1]⁻.

B. 2-(4-Fluorobenzenesulfonyl)benzene-1,4-diol

A solution of the title A compound, 4-fluorobenzenesulfinic acid (3 g, 18.75 mmol) in 10 mL of water is added to a solution of 1,4 benzoquinone (1.93 g, 17.86 mmol) in 30 mL of CH_2Cl_2 at RT. After 4 h, the precipitated product is collected by vacuum filtration and washed with cold CH_2Cl_2 and dried under vacuum to afford 2-(4-fluorobenzenesulfonyl)-benzene-1,4-diol: NMR (DMSO-d_6) 6.73 (d, 1H, $J = 9$), 6.92 (dd, 1H, $J = 9, 3$), 7.31 (d, 1H, $J = 3$), 7.41 (app t, 2H, $J = 9$), 7.96 (dd, 2H, $J = 9, 5$), 9.41 (s, 1H), 10.05 (s, 1H); ESI-MS 267 [M-1].

C. 4-(2,6-Dimethyl-4-nitrophenoxy)-2-(4-fluorobenzenesulfonyl)phenol

The title B compound, 2-(4-fluorobenzenesulfonyl)benzene-1,4-diol (1.2 g, 4.48 mmol) is added to a suspension of NaH (60% dispersion in mineral oil; 0.39 g, 9.86 mmol) in 15 mL of NMP at 0°C in one portion. The mixture is warmed to RT and after 30 min, 4-chloro-3,5-dimethylnitrobenzene (1 g, 5.38 mmol) is added and the reaction is heated at 120°C for 1 h. The reaction is cooled to RT and quenched with aqueous 1N HCl. The mixture is partitioned between water and EtOAc, and the organic solution is washed with water and brine, dried over anhydrous Na_2SO_4 and concentrated. Chromatography on silica (eluant; EtOAc/hexane - 1/2 / 1/1) affords 4-(2,6-dimethyl-4-nitrophenoxy)-2-(4-fluorobenzenesulfonyl)phenol: NMR (CDCl_3) 2.11 (s, 6H), 6.85- 6.96 (m, 3H), 7.00 (app t, 2H, $J = 9$), 7.88 (dd, 2H, $J = 9, 5$), 7.98 (s, 2H), 8.73 (s, 1H).

D. 4-(4-Amino-2,6-dimethylphenoxy)-2-(4-fluorobenzenesulfonyl)phenol

A mixture of the title C compound, 4-(2,6-dimethyl-4-nitrophenoxy)-2-(4-fluorobenzenesulfonyl)phenol (0.69 g, 1.65 mmol) and palladium on activated carbon (10 wt. %; 69 mg) in 10 mL of EtOH and 10 mL of CH_2Cl_2 is stirred under hydrogen atmosphere (H_2 , 1 atm) for 6 h. The catalyst is removed by vacuum filtration through celite, washed with a 1/1-mixture of EtOH and CH_2Cl_2 , and the combined filtrate and washings are concentrated and dried under vacuum to give 4-(4-amino-2,6-dimethylphenoxy)-2-(4-fluorobenzenesulfonyl)phenol: NMR (DMSO-d_6) 1.93 (s, 6H), 4.92 (s, 2H), 6.34 (s, 2H), 6.85 (d, 1H, $J = 9$), 7.00 (dd, 1H, $J = 9, 3$), 7.12 (d, 1H, $J = 3$), 7.43 (app t, 2H, $J = 9$), 7.94 (dd, 2H, $J = 9, 5$), 9.41 (s, 1H), 10.4 (s, 1H).

F002260-0969960


E. N-{4-[3-(4-Fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}oxamic acid ethyl ester

A mixture of the title D compound, 4-(4-amino-2,6-dimethylphenoxy)-2-(4-fluorobenzenesulfonyl)phenol (0.64 g, 1.65 mmol) and 2 mL of diethyl oxalate is heated at 180°C for 3 h. The reaction is cooled to RT and diethyl oxalate is removed under vacuum. Chromatography on silica (eluant; EtOAc/hexane - 1/3 → 2/3) affords N-{4-[3-(4-fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}oxamic acid ethyl ester: NMR (CDCl₃) 1.46 (t, 3H, J = 7.5), 2.06 (s, 6H), 4.42 (q, 2H, J = 7.5), 6.90-6.98 (m, 3H), 7.22 (app t, 2H, J = 8.3), 7.40 (s, 2H), 7.87-7.93 (m, 2H), 8.87 (br s, 1H); ESI-MS 486 [M-1].

F. N-{4-[3-(4-Fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}oxamic acid

A solution of the title E compound, N-{4-[3-(4-fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}oxamic acid ethyl ester (773 mg, 1.58 mmol) in 15 mL of EtOH is treated with aqueous 1N sodium hydroxide (NaOH; 4.75 mL, 4.75 mmol) at RT. After 1h, the reaction is quenched with aqueous 1N HCl (5.5 mL) and the product is taken up in EtOAc, washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The product is triturated with Et₂O and dried in vacuo to afford N-{4-[3-(4-fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}oxamic acid: NMR (DMSO-d₆) 2.06 (s, 6H), 6.88 (d, 1H, J = 9), 7.03 (dd, 1H, J = 9,3), 7.13 (d, 1H, J = 3), 7.43 (app t, 2H, J = 9), 7.61 (s, 2H), 7.94 (dd, 2H, J = 9,5), 10.5 (s, 1H), 10.69 (s, 1H); IR (KBr) 1240, 1481, 1685, 1764; ESI-MS 458 [M-1].

The following additional compounds are similarly prepared.




Example	NMR	IR (KBr)	ESI-MS
R4			
27 R4 = Ph-	(MeOH-d ₆): 2.02 (s, 6H), 6.85 (d, 1H, J = 7), 6.98 (dd, 1H, J = 7, 2.3), 7.26 (d, 1H, J = 2.3), 7.53 (s, 2H), 7.55-7.68 (m, 3H), 7.93 (d, 2H, J = 7)	1209, 1479, 1697	440 [M-1] ⁻
28 R4 = p-ClC ₆ H ₄ -	(MeOH-d ₆): 2.12 (s, 6H), 6.84 (d, 1H, J = 8.3), 6.96 (dd, 1H, J = 8.3, 3), 7.24 (d, 1H, J = 3), 7.53 (d, 2H, J = 7.5), 7.58 (s, 2H), 7.90 (d, 2H, J = 7.5)	1207, 1479, 1697	474 [M-1] ⁻
29 R4 = p-MeC ₆ H ₄ -	(MeOH-d ₆): 2.10 (s, 6H), 2.41 (s, 3H), 6.82 (d, 1H, J = 9), 6.95 (dd, 1H, J = 9, 3), 7.19 (d, 1H, J = 3), 7.45 (d, 2H, J = 8.3), 7.53 (s, 2H), 7.77 (d, 2H, J = 8.3)	1149, 1207, 1481	454 [M-1] ⁻ 473 [M+NH ₄] ⁺
30 R4 = p-MeOC ₆ H ₄ -	(CDCl ₃): 1.98 (s, 6H), 3.80 (s, 3H), 6.80-6.82 (m, 2H), 6.89-6.93 (m, 3H), 7.31 (s, 2H), 7.73 (d, 2H, J = 7.2), 8.73 (br s, 1H), 8.89 (br s, 1H)	1209, 1475, 1588, 1705	470 [M-1] ⁻
31 R4 = p-CF ₃ C ₆ H ₄ -	(DMSO-d ₆): 2.07 (s, 6H), 6.89 (d, 1H, J = 9), 7.08 (dd, 1H, J = 9, 3), 7.15 (d, 1H, J = 3), 7.60 (s, 2H), 7.98 (d, 2H, J = 3), 8.08 (d, 2H, J = 3)	1207, 1476, 1697	508 [M-1] ⁻
32 R4 = Me-	(MeOH-d ₄): 2.12 (s, 6H), 3.23 (s, 3H), 6.96-7.0 (m, 2H), 7.11 (d, 1H, J = 2.2), 7.51 (s, 2H)	1205, 1481, 1687, 1739	378 [M-1] ⁻

T09260" 09699660

33 R4 = n-Bu-	(DMSO-d ₆): 0.81 (t, 3H, J = 7.2), 1.24-1.36 (m, 2H), 1.41-1.51 (m, 2H), 2.04 (s, 6H), 3.37 (t, 2H, J = 7.2), 6.88 (d, 1H, J = 3), 7.08 (dd, 1H, J = 9, 3), 7.15 (d, 1H, J = 3), 7.00-7.10 (m, 2H), 7.57 (s, 2H), 10.67 (s, 2H)	1205, 1488, 1694	420 [M-1] ⁻ 439 [M+NH ₄] ⁺
34 R4 = i-Pr-	(DMSO-d ₆): 1.11 (d, 6H, J = 7), 2.03 (s, 6H), 3.66-3.75 (m, 1H), 6.88 (d, 1H, J = 3), 7.02 (d, 1H, J = 3), 7.08 (dd, 1H, J = 9, 3), 7.57 (s, 2H), 10.65 (s, 2H)	1229, 1354, 1480, 1688, 1761	406 [M-1] ⁻ 425 [M+NH ₄] ⁺

Example 35


N-{4-[3-(4-Fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}malonic acid



The title compound is prepared similarly to Example 26: NMR (DMSO-d₆) 2.05 (s, 6H), 3.35 (s, 2H), 6.89 (d, 1H, J = 9), 7.01 (dd, 1H, J = 9, 3), 7.14 (d, 1H, J = 3), 7.39-7.45 (m, 4H), 7.93 (dd, 2H, J = 8.8, 5.2), 10.18 (s, 1H), 10.55 (s, 1H), 12.6 (br s, 1H); IR (KBr) 1142, 1239, 1485, 1623, 1654, 1736; ESI-MS 472 [M-1]⁻.

Example 36

N-{4-[3-(4-Fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}succinamic acid




The title compound is prepared similarly to Example 26: NMR (DMSO-d₆) 2.04 (s, 6H), 2.50-2.56 (m, 4H), 6.87 (d, 1H, J = 9), 7.01 (dd, 1H, J = 9, 3), 7.12 (d, 1H, J = 3), 7.39-7.45 (m,

T02260-09699660

4H), 7.93 (dd, 2H, J = 8.8, 5.2), 9.93 (s, 1H), 10.5 (br s, 1H), 12.1 (s, 1H); IR (KBr) 1480, 1659, 1717; ESI-MS 486 [M-1]⁻.

Example 37


3-{4-[3-(4-Fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenylamino}-propionic acid



The title compound is prepared similarly to Example 26: NMR (MeOH-d₄) 2.06 (s, 6H), 2.65 (t, 2H, J = 7), 3.49 (t, 2H, J = 7), 6.82 (d, 1H, J = 9), 6.85 (s, 2H), 6.94 (dd, 1H, J = 9, 3), 7.15 (d, 1H, J = 3), 7.23 (app t, 2H, J = 9), 7.88-7.93 (m, 2H); IR (KBr) 1199, 1493, 1675; ESI-MS 460 [M+1]⁺, 458 [M-1]⁻.

Example 38


N-{4-[3-(4-Fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3-methylphenyl}oxamic acid



The title compound is prepared similarly to Example 26: NMR (DMSO-d₆) 2.18 (s, 3H), 6.90 (d, 1H, J = 9), 6.92 (d, 1H, J = 9), 7.16 (dd, 1H, J = 9, 3), 7.34 (d, 1H, J = 3), 7.43 (app t, 2H, J = 9), 7.61 (dd, 1H, J = 9, 3), 7.74 (d, 1H, J = 3), 7.97 (dd, 2H, J = 9, 5), 10.67 (s, 1H), 10.71 (s, 1H); IR (KBr) 1234, 1495, 1697; ESI-MS 444 [M-1]⁻.

Example 39


N-{3,5-Dibromo-4-[3-(4-fluorobenzenesulfonyl)-4-hydroxyphenoxy]phenyl}oxamic acid



SECRET

SECRET


SECRET



SECRET

SECRET


SECRET



SECRET

Example 42


N-{4-[3-(4-Fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}-N'-isopropyl-oxalamide



The title compound is prepared analogously to Example 26: ES-MS 499 [M-1]⁻, 518 [M+NH₄]⁺.

Example 43


N-Butyl-N'-{4-[3-(4-fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}-oxalamide



The title compound is prepared analogously to Example 26: ES-MS 513 [M-1]⁻, 515 [M+1]⁺, 532 [M+NH₄]⁺.

Example 44

N-{4-[3-(4-Fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}-N'-(2-methoxyethyl)oxalamide




The title compound is prepared analogously to Example 26: ES-MS 517 [M+1]⁺.

0966960-092604

Example 45


N-{4-[3-(4-Fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}-2-morpholin-4-yl-2-oxoacetamide



The title compound is prepared analogously to Example 26: ES-MS 527 $[M-1]^-$, 529 $[M+1]^+$, 546 $[M+NH_4]^+$.

Example 46

N-{4-[4-Hydroxy-3-(piperidine-1-carbonyl)phenoxy]-3,5-dimethylphenyl}oxamic acid



A. 5-(2,6-Dimethyl-4-nitrophenoxy)-2-(2-methoxyethoxymethoxy)benzoic acid 2-methoxyethoxymethyl ester

A suspension of sodium NaH (60% dispersion in mineral oil; 1.32 g, 33 mmol) in 50 mL of NMP is cooled to 0°C and 2,5-dihydroxybenzoic acid (1.54 g, 10 mmol) is added in one portion. The mixture is warmed to room temperature, and after 30 min, 4-chloro-3,5-dimethylnitrobenzene (2.41 g, 13 mmol) is added in one portion and the reaction is heated at 120°C for 3 h. The reaction is cooled to RT and 2-methoxyethoxymethyl chloride (2.85 mL, 25 mmol) is added. After stirring for 30 min, the mixture is poured onto water and the product is taken up in Et₂O. The organic solution is washed with brine, dried over anhydrous Na₂SO₄ and concentrated. Chromatography on silica (eluant; EtOAc/hexane - 1/2 → 3/2) affords 5-(2,6-dimethyl-4-nitrophenoxy)-2-(2-methoxyethoxymethoxy)benzoic acid 2-methoxyethoxymethyl ester: NMR (CDCl₃) 2.23 (s, 6H), 3.36 (s, 3H), 3.38 (s, 3H), 3.54-3.60 (m, 4H), 3.80-3.90 (m,

F022610-09699560

4H), 5.28 (s, 2H), 5.52 (s, 2H), 6.82 (dd, 1H, J = 9,3), 7.17-7.23 (m, 2H), 8.02 (s, 2H).

B. 5-(4-Amino-2,6-dimethylphenoxy)-2-(2-methoxyethoxymethoxy)benzoic acid 2-methoxyethoxymethyl ester

A mixture of the title A compound, 5-(2,6-dimethyl-4-nitrophenoxy)-2-(2-methoxyethoxymethoxy)benzoic acid 2-methoxyethoxymethyl ester (3.2 g, 6.68 mmol) and palladium on activated carbon (10 wt. %; 320 mg) in 50 mL of EtOAc is stirred under hydrogen atmosphere (H₂, 1 atm) for 3 h. The catalyst is removed by vacuum filtration through celite, washed with EtOAc, and the combined filtrate and washings are concentrated and dried under vacuum to give 5-(4-amino-2,6-dimethylphenoxy)-2-(2-methoxyethoxymethoxy)benzoic acid 2-methoxyethoxymethyl ester: NMR (CDCl₃) 2.07 (s, 6H), 3.37 (s, 3H), 3.40 (s, 3H), 3.52-3.62 (m, 4H), 3.83-3.92 (m, 4H), 5.26 (s, 2H), 5.52 (s, 2H), 6.60 (s, 2H), 6.80 (dd, 1H, J = 8.3, 3), 7.13 (d, 1H, J = 8.3), 7.24 (d, 1H, J = 3).

C. 5-[4-(Ethoxyoxalylamino)-2,6-dimethylphenoxy]-2-hydroxybenzoic acid

A solution of the title B compound, 5-(4-amino-2,6-dimethylphenoxy)-2-(2-methoxyethoxymethoxy)benzoic acid 2-methoxyethoxymethyl ester (2.83 g, 6.3 mmol) in 20 mL of THF is cooled to 0°C and treated sequentially with NMM (2.1 mL, 18.9 mmol) and ethyl oxalyl chloride (0.915 mL, 8.19 mmol). After 15 min, the mixture is partitioned between EtOAc and water, and the organic solution is washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The residue is dissolved in 30 mL of EtOH and 20 mL of aqueous 6N HCl is added. The mixture is stirred at RT for 16 h, and EtOH is removed under reduced pressure. The residue is diluted with water (100 mL), and the solid is collected by vacuum filtration, washed with water and dried. Crystallization from acetonitrile gives 5-[4-(ethoxyoxalylamino)-2,6-dimethylphenoxy]-2-hydroxybenzoic acid: NMR (DMSO-d₆) 1.31 (t, 3H, J = 7), 2.06 (s, 6H), 4.30 (q, 2H, J = 7), 6.88-6.97 (m, 2H), 7.08 (dd, 1H, J = 9, 3), 7.54 (s, 2H), 10.7 (s, 1H).


D. N-{4-[4-Hydroxy-3-(piperidine-1-carbonyl)phenoxy]-3,5-dimethylphenyl}oxamic acid

0966660-092301

A solution of title C compound, 5-[4-(ethoxyoxalylamino)-2,6-dimethylphenoxy]-2-hydroxybenzoic acid (37 mg, 0.1 mmol) in 1 mL of DMF is treated with NMM (55 μ L, 0.5 mmol) and 1,1'-carbonyldiimidazole (32 mg, 0.2 mmol) at RT. The reaction mixture is heated at 60°C for 1 h, then cooled to RT and piperidine (24 μ L, 0.24 mmol) is added. After 16 h, the reaction is treated with aqueous 1.5 N LiOH (333 μ L, 0.5 mmol). The mixture is agitated for 30 min, and the reaction is quenched with trifluoroacetic acid (TFA; 100 μ L). The product is purified by HPLC (mobile phase; acetonitrile-water with 0.1% of trifluoroacetic acid) to afford N-{4-[4-hydroxy-3-(piperidine-1-carbonyl)phenoxy]-3,5-dimethylphenyl}oxamic acid: NMR (DMSO- d_6) 1.34-1.63 (m, 6H), 2.06 (s, 6H), 3.30 (br s, 4H), 6.36 (d, 1H, J = 2.3), 6.67 (dd, 1H, J = 8.3, 2.3), 6.80 (d, 1H, J = 8.3), 7.53 (s, 2H), 9.38 (br s, 1H), 10.6 (s, 1H); ESI-MS 413 [M+1]⁺.

Example 47


N-{4-[4-Hydroxy-3-(morpholine-4-carbonyl)phenoxy]-3,5-dimethylphenyl}oxamic acid



The title compound is prepared similarly to Example 46: ESI-MS 415 [M+1]⁺.

Example 48

N-[4-(3-Cyclohexylcarbonyl-4-hydroxyphenoxy)-3,5-dimethylphenyl]oxamic acid




The title compound is prepared similarly to Example 46: ESI-MS 427 [M+1]⁺.

T092250*09099660

Example 49


N-{4-[4-Hydroxy-3-(2-methoxyethylcarbamoyl)phenoxy]-3,5-dimethylphenyl}oxamic acid



The title compound is prepared similarly to Example 46: ESI-MS 403 [M+1]⁺.

Example 50


N-{4-[4-Hydroxy-3-(2-morpholin-4-yl-ethylcarbamoyl)phenoxy]-3,5-dimethylphenyl}-oxamic acid



The title compound is prepared similarly to Example 46: ESI-MS 458 [M+1]⁺.

Example 51

N-{4-[4-Hydroxy-3-(pyridin-3-ylcarbamoyl)phenoxy]-3,5-dimethylphenyl}oxamic acid



The title compound is prepared similarly to Example 46: ESI-MS 422 [M+1]⁺.

T02250-09699660

Example 52

The following examples can be prepared analogously as described in the previous Examples:

N-[3,5-Dichloro-4-(3-cyclopentanesulfonyl-4-hydroxyphenoxy)phenyl]oxamic acid;
N-[3,5-Dichloro-4-(3-cyclopropylmethanesulfonyl-4-hydroxyphenoxy)-phenyl]oxamic acid;
N-[3,5-Dichloro-4-(3-cyclobutylmethanesulfonyl-4-hydroxyphenoxy)phenyl]oxamic acid;
N-[4-(3-Cyclopropylmethanesulfonyl-4-hydroxyphenoxy)-3,5-dimethylphenyl]oxamic acid;
N-[3-Chloro-4-(3-cyclobutylmethanesulfonyl-4-hydroxyphenoxy)-5-methylphenyl]oxamic acid;
N-[4-(3-Cyclobutylmethanesulfonyl-4-hydroxyphenoxy)-3,5-dimethylphenyl]oxamic acid;
N-[4-(3-Cyclopentylmethanesulfonyl-4-hydroxyphenoxy)-3,5-dimethylphenyl]oxamic acid;
N-[3-Chloro-4-(3-cyclopentylmethanesulfonyl-4-hydroxyphenoxy)-5-methylphenyl]oxamic acid;
N-[3,5-Dichloro-4-(3-cyclopentylmethanesulfonyl-4-hydroxyphenoxy)phenyl]oxamic acid;
N-[4-(3-Cyclohexylmethanesulfonyl-4-hydroxyphenoxy)-3,5-dimethylphenyl]oxamic acid;
N-[3-Chloro-4-(3-cyclohexylmethanesulfonyl-4-hydroxyphenoxy)-5-methylphenyl]oxamic acid;
N-[3,5-Dichloro-4-(3-cyclohexylmethanesulfonyl-4-hydroxyphenoxy)phenyl]oxamic acid;
N-[3,5-Dichloro-4-[3-(4-fluorobenzenesulfonyl)-4-hydroxyphenoxy]phenyl]oxamic acid;
N-[3-Chloro-4-[3-(4-fluorobenzenesulfonyl)-4-hydroxyphenoxy]-5-methylphenyl]oxamic acid;
N-[3,5-Dichloro-4-[3-(4-chlorobenzenesulfonyl)-4-hydroxyphenoxy]-phenyl]oxamic acid ethyl ester;
N-[3,5-Dichloro-4-[3-(4-chlorobenzenesulfonyl)-4-hydroxyphenoxy]phenyl]oxamic acid;
N-[3-Chloro-4-[3-(4-chlorobenzenesulfonyl)-4-hydroxyphenoxy]-5-methylphenyl]oxamic acid;
N-[3-Chloro-4-[3-(4-chlorobenzenesulfonyl)-4-hydroxyphenoxy]-5-methylphenyl]oxamide;
N-[4-(3-Benzenesulfonyl-4-hydroxyphenoxy)-3,5-dichlorophenyl]oxamic acid ethyl ester;
N-[4-(3-Benzenesulfonyl-4-hydroxyphenoxy)-3,5-dichlorophenyl]oxamic acid;
N-[4-(3-Benzenesulfonyl-4-hydroxyphenoxy)-3,5-dichlorophenyl]oxamide;

T002269-090900

N-{3,5-Dichloro-4-[4-hydroxy-3-(naphthalene-1-sulfonyl)phenoxy]phenyl}oxamic acid ethyl ester;
N-{3,5-Dichloro-4-[4-hydroxy-3-(naphthalene-1-sulfonyl)phenoxy]phenyl}oxamic acid;
N-{3,5-Dichloro-4-[4-hydroxy-3-(naphthalene-2-sulfonyl)phenoxy]phenyl}oxamic acid ethyl ester;
N-{3,5-Dichloro-4-[4-hydroxy-3-(naphthalene-2-sulfonyl)phenoxy]phenyl}oxamic acid;
N-{3,5-Dichloro-4-[4-hydroxy-3-(toluene-4-sulfonyl)phenoxy]phenyl}oxamic acid;
N-{3,5-Dichloro-4-[4-hydroxy-3-(toluene-4-sulfonyl)phenoxy]phenyl}oxamide;
N-[3,5-Dichloro-4-(3-cyclopropylmethanesulfonyl-4-hydroxyphenoxy)phenyl]oxamic acid ethyl ester;
N-[4-(4-Hydroxy-3-methanesulfonylphenoxy)-3,5-dimethylphenyl]oxamic acid;
N-[3,5-Dichloro-4-(3-ethanesulfonyl-4-hydroxyphenoxy)phenyl]oxamic acid;
N-{3,5-Dichloro-4-[4-hydroxy-3-(propane-2-sulfonyl)phenoxy]phenyl}oxamic acid ethyl ester;
N-[3,5-Dichloro-4-(3-cyclobutylmethanesulfonyl-4-hydroxyphenoxy)phenyl]oxamic acid ethyl ester;
N-[3,5-Dichloro-4-(3-cyclohexylmethanesulfonyl-4-hydroxyphenoxy)phenyl]oxamic acid ethyl ester;
N-[3,5-Dichloro-4-(3-cyclopentanesulfonyl-4-hydroxyphenoxy)phenyl]oxamic acid ethyl ester;
N-[4-[3-(Butane-1-sulfonyl)-4-hydroxyphenoxy]-3,5-dichlorophenyl]oxamic acid;
N-[3,5-Dichloro-4-(4-hydroxy-3-phenylmethanesulfonylphenoxy)phenyl]oxamic acid;
N-{3,5-Dichloro-4-[4-hydroxy-3-(propane-1-sulfonyl)phenoxy]phenyl}oxamic acid;
N-{3,5-Dichloro-4-[3-(4-fluorophenylmethanesulfonyl)-4-hydroxyphenoxy]phenyl}oxamic acid.

Example 53

The following examples can be prepared as described herein in the specification or according to methods known in the art:

1H-Tetrazole-5-carboxylic acid {4-[3-(4-fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}amide;

1H-Tetrazole-5-carboxylic acid {3-chloro-4-[3-(4-fluorobenzenesulfonyl)-4-hydroxyphenoxy]-5-methylphenyl}amide;

1H-Tetrazole-5-carboxylic acid {3,5-dichloro-4-[3-(4-fluorobenzenesulfonyl)-4-hydroxyphenoxy]-phenyl}amide.

Example 54

Formulation Example A: Hard gelatin capsules, comprising 100 mg active substance can be prepared for example as follows:

Composition (for 1000 capsules)

Active ingredient	100.0 g
Lactose	250.0 g
Microcrystalline cellulose	30.0 g
Sodium lauryl sulfate	2.0 g
Magnesium stearate	8.0 g

The sodium lauryl sulfate is added to the lyophilized active ingredient via a sieve with a mesh size of 0.2 mm. Both components are intimately mixed. Then first the lactose is added via a sieve with a mesh size of 0.6 mm and then the microcrystalline cellulose via a sieve with a mesh size of 0.9 mm. Thereupon these components are intimately mixed for a further 10 minutes. Finally the magnesium stearate is added via a sieve with a mesh size of 0.8 mm. After 3 minutes of further mixing, 390 mg each of the formulation obtained are filled into hard gelatin capsules of size 0.

Formulation Example B: Hard gelatin capsules comprising either 0.1 or 0.3 mg of active ingredient, N-{4-[3-(4-fluorobenzenesulfonyl)-4-hydroxyphenoxy]-3,5-dimethylphenyl}oxamic acid, can be prepared analogously as described in Formulation example A:

Composition (for one capsule)

Active ingredient	0.100 mg	0.300 mg
Microcrystalline cellulose	73.775 mg	73.575 mg
Sodium lauryl sulfate	0.750 mg	0.750 mg
Magnesium stearate	0.375 mg	0.375 mg

108260-0969660